

Taxonomic studies in *Thomasia* and *Lasiopetalum* (Sterculiaceae)

by Susan Paust*

Abstract

Four new species of *Thomasia* and six new species of *Lasiopetalum* are described; *Thomasia formosa* sp. nov., *T. gardneri* sp. nov., *T. microphylla* sp. nov., *T. tremandroides* sp. nov.; *Lasiopetalum cardiophyllum* sp. nov., *L. compactum* sp. nov., *L. glabratum* sp. nov., *L. lineare* sp. nov., *L. monticola* sp. nov. and *L. rotundifolium* sp. nov.

Two species of *Lasiopetalum* are reduced to synonymy: *L. angustifolium* W. V. Fitzg. is synonymous with *L. oppositifolium* F. Muell.; and *L. acutiflorum* Turcz. with *L. indutum* Steud.

All specimens cited are housed at PERTH, except where otherwise indicated.

Thomasia tremandroides S. Paust sp. nov. (Figures 1 and 11) (tremandroides = resembling some species of Tremandraceae)

Folia ovata, brevissime petiolata, 4–8 x 2·5–5 mm stellato-tomentosa. *Stipulae* foliaceae, 2·5 mm longae. *Racemus* pilis stellatis et simplici-moniliformibus tomentosus. *Braeteolae* 3, anguste lineares, calycem subtus. *Calyx* 8 mm longus, stellato-hirsutus lobis 5 ovatis. *Petala* 5, spatulata, 1 mm longa. *Antherae* 5, sere sessiles, anguste triangulares, 3 mm longae. *Staminodia* 5, clavata, 1 mm longa. *Ovarium* 3-loculare, 1 mm longum pilis glandulosis et stellatis sparsim tomentosum. *Style* 2 mm longus, glaber.

Type: 33 mi N of Wubin towards Percnjori, Western Australia, 8 Oct. 1972, S. Paust 1319. Holo: PERTH; iso: CANB, K.

Erect, multistemmed, woody perennial, 10–60 cm high. Branchlets tomentose with stellate and simple-moniliform hairs. Leaves alternate; petiole minute; lamina ovate, chartaceous, 4–8 mm long, 2·5–5 mm wide, stellate-tomentose; margin undulate, slightly recurved. Stipules foliaceous, ovate-orbicular, often oblique, 2·5 mm long. Racemes leaf-opposed, 20–40 mm long, 1–4 flowered, indumentum similar to that of the branchlets; peduncle 10–30 mm long; pedicel 3–7 mm long; bracts linear, 2 mm long; bracteoles 3, narrow linear, 5 mm long, subtending calyx. Calyx angular in bud, mauve-pink, exuplicate and papery at anthesis, 8 mm long, stellate-hairy, divided to below the middle into 5 ovate lobes, ribs thickened especially at base. Petals 5, spatulate, 1 mm long, purple-black. Anthers 5, almost sessile, narrowly triangular, 3 mm long, purple-black. Staminodes 5, obscured by anthers, clavate, 1 mm long, alternate and basally connate with stamens. Ovary globose, 1 mm long, purple-black, with scattered simple, glandular and stellate hairs, 3-celled; style filiform, 2 mm long, glabrous, yellowish; ovules 3–4 per locule. Seed not seen.

Distribution: South-west Western Australia; from Bencubbin north to the Murchison River.

Murchison River, Oct. 1961, de la Hunt; Boolardy Station, A. B. Cashmore 128; Canna, C. A. Gardner 2691; near Morowa, W. E. Blackall 2817; 1 mi S of Caron, S. Paust 1309; 9·2 mi E of Bindy Bindy, S. Paust 1026; 6 mi S of Ballidu, J. W. Green 791; 13 mi N of Bencubbin, W. E. Blackall 3317; (?) Mt. Barker to Walpole, 31 Aug. 1965, F. W. Humphreys. (It is extremely doubtful that the collection data of the Humphreys specimen are correct.)

Thomasia tremandroides is distinguished from other species of *Thomasia* by its small papery ovate leaves and by the presence and morphology of its staminodes.

* Present address: Mrs. R. Downes, Cleator, Cumberland, England.



Western Australian Herbarium

Thomasia tremandrodes S. Paust

DETERMINAVIT Susan Paust June 1973

WESTERN AUSTRALIAN HERBARIUM, PERTH
Flora of Western Australia

Thomasia

Calyx pink-purple, base darker becoming cream coloured at base.
Petals and stamens deep purple-black. Stems 2 ft tall branched
nearly together, sparsely yellowish pubescent.

Loc. 58 mi N of Albany towards Kerenjou

Coll. Susan Paust 1319 8/1/1972

44-14-33 00000000

WESTERN AUSTRALIAN HERBARIUM, PERTH
Flora of Western Australia

Loc.

Coll.

19

Figure 1—*Thomasia tremandrodes* sp. nov. Holotype—S. Paust 1319.



Figure 2—*Thomasia formosa* sp. nov. Holotype—± 5 mi W of Three Springs, C. Chapman.

Thomasia formosa S. Paust sp. nov. (Figures 2 and 11)
(formosa = handsome)

Frutex stellato-tomentosus. *Folia* angustissime ovata, petiolata, 20–40 x 4–10 mm. *Stipulae* oblique ovatae, 4–7 mm longae. *Racemi* saepe ramosi. *Bracteolae* 3, lanceolatae, calycem subtus. *Calyx* 10 mm longus, stellato-hirsutus, lobis 5 ovatis. *Petala* 5, spatulata, 1 mm longa. *Stamina* 5, filamentis 1 mm longis, antheris anguste triangularibus 3·5 mm longis. *Ovarium* 3-loculare, 2 mm longum, albo-stellato-tomentosum. *Stylus* 3 mm longus, glaber.

Type: ± 5 mi W of Three Springs, Western Australia, 29 Sept. 1972. C. Chapman s.n
Holo: PERTH, iso: K.

Erect, multistemmed, woody *perennial*, up to 50 cm high; branchlets, leaves and inflorescence closely stellate-tomentose, rufous. Leaves alternate; petiole 5 mm long; lamina very narrowly ovate, 20–40 mm long, 4–10 mm wide, rugose above; margin ± crenulate, recurved. Stipules foliaceous, obliquely ovate, 4–7 mm long. Racemes leaf-opposed, 70–120 mm long, 7- many-flowered, often branched; peduncle 30–40 mm long; pedicels 10 mm long; bracts ovate 1–3 mm long; bracteoles 3, lanceolate, reflexed, 7 mm long, subtending calyx. Calyx pink, papery, 10 mm long, stellate-hairy, divided to below the middle into 5 ovate lobes, ribs thickened. Petals 5, spatulate, slightly concave, 1 mm long, glabrous. Stamens 5; filaments linear, 1 mm long; anthers narrowly triangular, 3·5 mm long. Ovary globose, 2 mm long, white stellate-tomentose, 3-celled; style filiform, 3 mm long, glabrous, ovules 4–8 per locule. Seed not seen.

Distribution: South-west Western Australia; this species is known only from collections made 8 km W of Three Springs.

Thomasia formosa is similar to the more southern species, *T. angustifolia* Steud. and *T. petalocalyx* F. Muell. in that it has a tomentose, 3-celled ovary and a glabrous style. It is distinguished from these species by the much larger flowers and inflorescence and by its leaf form.

Thomasia microphylla S. Paust. sp. nov. (Figures 3 and 11)
(micro = small, phyllum = leaf)

Frutex squamis ut videtur ciliatis lepidotus. *Folia* ovato-oblonga, breviter petiolata, 2–10 x 1·5–5 mm, coriacea, supra glabra, subtus lepidota. *Stipulae* nullae. *Racemi* 1–2-floribus. *Bracetolae* 3, teretes. *Calyx* 8 mm longus, varie-lepidotus, lobis 5 ovatis. *Petala* 5, ovata, carnosa, 0·8 mm longa. *Stamina* 5, filamentis 1 mm longis, antheris oblongo-linearibus, 3 mm longis. *Staminodia* ± 5, teretia, 1·5 mm longa. *Ovarium* 3-loculare, 2 mm longum, dense lepidotum. *Stylus* 3 mm longus, praeter basem glaber.

Type: Mt. Short, at N end of Ravensthorpe Range, Western Australia, 30 Aug. 1963, A. S. George 5698. Holo: PERTH, iso: CANB, K.

Erect, multistemmed, woody *perennial*, 15–35 cm high, branchlets, leaves and inflorescence variably lepidote with apparently ciliate scales. Leaves alternate, slightly reflexed; petiole 1–2 mm long, lepidote; lamina ovate-oblong, slightly cordate, 2–10 mm long, 1·5–5 mm wide, entire, coriaceous, smooth to reticulate and glabrous above, lepidote below. Stipules absent. Racemes terminal, 10–40 mm long, 1–2 flowered, peduncle 10–30 mm long; pedicels 4–5 mm long; bracts terete, 1–1·5 mm long, often absent; bracteoles 3, terete, 1–2 mm long, subtending calyx. Calyx angular in bud, pale mauve, 7–9 mm long, finely stellate-lepidote, the outer surface with additional large brown scales, particularly on the prominent ribs, divided almost to the base into 5 broadly ovate lobes. Petals 5, ovate, concave, fleshy, normally glabrous, 0·8 mm long. Stamens 5; filaments oblong, 1 mm long; anthers oblong-linear, 3 mm long, shortly attenuate and yellow at apex. Staminodes 5, alternate with stamens, terete, 1·5 mm long, often absent. Ovary ovoid, 2 mm long, densely lepidote, 3-celled; style filiform, 3 mm long, sparsely lepidote at base, otherwise glabrous; ovules 2–3 per locule. Fruit a loculicidal capsule; seed 1 per locule, almost cylindrical, villous, 2 mm long, 0·9 mm wide.



Thomasia microphylla S. Paust

DETERMINAVIT Susan Paust June 1973

WESTERN AUSTRALIAN HERBARIUM
PERTH WA

Much branched, spreading shrub to 50cm.
Fls pink to bluish, among thick
mallee scrub.

St. Lant. St V end of Ravensthorpe
Loc. Lang. 11
Coll. A. S. GEORGE 5691 20 Aug 1963

1963

WESTERN AUSTRALIAN HERBARIUM, PERTH
Flora of Western Australia

Loc.
Coll.

19

Figure 3—*Thomasia microphylla* sp. nov. Holotype—A. S. George 5698.

Distribution: South-west Western Australia; from Ongerup to east of Esperance.

10 mi E of Ongerup, K. Newbey 379; 22 mi S of Ravensthorpe, E. Wittwer 397; 18 mi E of Ravensthorpe, K. Newbey 1312; 40 mi E and 10 mi N of Esperance turn off, F. Lullfitz 3579.

Thomasia microphylla has affinities with *T. stelligera* (Turez.) Benth., but is readily distinguished from that species by its smaller, ovate-oblong leaves.

***Thomasia gardneri* S. Paust sp. nov. (Figures 4 and 11)**

(named after the late C. A. Gardner, the collector of the type specimen and former Government Botanist of Western Australia)

Frutex squamis ut videtur ciliatis lepidotus. *Folia* anguste-ovata, breviter petiolata, 8–20 x 4–8 mm, coriacea, supra glabra, subtus lepidota. *Stipulae* nullae. *Racemi* 1–2-floribus. *Bracteolae* 3, teretes. *Calyx* 9 mm longus, intus stellato-hirsutus, extus lepidotus, lobis 5 ovatis. *Petala* 5, spatulata, chartacea, 1·5 mm longa, marginibus sparsim lepidotis. *Stamina* 5, filamenta 2 mm longis, antheris oblongis, 2 mm longis. *Ovarium* 5-loculare, 3 mm longum, dense lepidotum. *Stylus* 5 mm longus, praeter basem glaber.

Type: Near Mt. Holland, Western Australia, Sept. 1929, C. A. Gardner s.n. Holo: PERTH.

Erect, multistemmed, woody *perennial*, up to 50 cm high; branchlets, leaves and inflorescences variably lepidote with apparently ciliate scales. *Leaves* alternate, slightly reflexed; petiole 2–4 mm long, lepidote; lamina narrowly ovate, 8–20 mm long, 4–8 mm wide, slightly cordate, entire, coriaceous, smooth to reticulate and glabrous above, lepidote below. *Stipules* absent. *Racemes* terminal, 10–30 mm long, 1–2 flowered; peduncle 5–15 mm long; pedicels slightly elevata, 5 mm long; bracts terete, often absent, 2 mm long; bracteoles 3, terete, 2 mm long, subtending calyx. *Calyx* prominently veined, pink, 9 mm long, sparsely stellate-hairy within, lepidote outside (especially on the ribs), divided almost to the base into 5 ovate lobes. *Petals* 5, spatulate, papery, 1·5 mm long, apical margin sparsely lepidote otherwise glabrous. *Stamens* 5, filaments oblong, 2 mm long; anthers oblong, 2 mm long. *Staminodes* absent. *Ovary* ovoid, 3 mm long, densely lepidote, 5-celled; style simple, 1·5 mm long, sparsely lepidote at base, otherwise glabrous; ovules 2 per locule. *Seed* not seen.

Distribution: South-west Western Australia; near Mt. Holland.

Thomasia gardneri is known only from the type and other collections from the same locality made by C. A. Gardner in September 1929. Its affinities lie with *T. microphylla* S. Paust (see above), but it is distinguished by its larger leaves, 5-celled ovary and differently-shaped petals and anthers.

***Lasiopetalum lineare* S. Paust. sp. nov. (Figures 5 and 12)**

(*linearis* = linear, referring to the leaves)

Folia anguste-linearia, brevissime petiolata, 20–40 x 1–4 mm, supra glabra, subtus stellato-tomentosa, marginibus revolutis. *Cymae* in capitula laxa contractae. *Bracteolae* filiformes. *Calyx* 7 mm longus, intus glaber, extus stellato-tomentosus, lobis 5, lineari-lanceolatis. *Petala* 5, orbicularia, 0·5 mm longa. *Antherae* fere sessiles, late-oblongae, 1·5 mm longae. *Ovarium* 3-loculare, 1 mm longum, albo-stellato-tomentosum. *Stylus* 3 mm longus pilis stellatis grandibus reflexis albis hirsutus.

Type: Watheroo West, Western Australia, 4 Nov. 1954, R. D. Royce 4965. Holo: PERTH, iso: K.

Erect, multistemmed, woody perennial, up to 40 cm high. *Branchlets* stellate-tomentose. *Leaves* alternate; petiole minute; lamina narrowly linear, 20–40 mm long, 1–4 mm wide, normally tightly revolute, almost glabrous above, stellate-tomentose below. *Cymes* leaf opposed, ± 8-flowered, closely stellate-tomentose, contracted into loose heads; peduncle 10–20 mm long; bracts filiform, 3 mm long; bracteoles normally 1, filiform, 4 mm long, subtending calyx. *Calyx* lilac (dark maroon near base of anthers), 7 mm long, almost glabrous within, white stellate tomentose outside, divided to the base into 5 linear-lanceolate lobes. *Petals* 5, orbicular, 0·5 mm long. *Anthers* 5, almost sessile,



HERBARIUM GARDNERIANUM

Made available to the

WESTERN AUSTRALIAN HERBARIUM

by the Lord Abbot and Members
of the Benedictine Community of
New Norcia June 1970

Spec. No. 100

C. A. Gardner: *Plantae Australiae occidentalis*

N.

1970-1971
Mt. Holland
Locality 1000 ft.
Flora of Western Australia

leg. C. A. Gardner

19

HOLOTYPE

Western Australian Herbarium
Thomasia gardneri S. Paust

TERMINATIVE Suan Paust June 1973




Figure 4—*Thomasia gardneri* sp. nov. Holotype—Near Mt. Holland, C. A. Gardner.



HOLOTYPE

W. Austr. Fl. Exs.

Lasiopetalum lineare S. Paust

DETERMINAVIT S. Paust June 1973

WESTERN AUSTRALIAN HERBARIUM, PERTH
Flora of Western Australia

STATE HERBARIUM
WESTERN AUSTRALIA

Loc.
Coll. R. D. ROYCE

19

Loc.
Coll.

19

Figure 5—*Lasiopetalum lineare* sp. nov. Holotype—R. D. Royce 4965.

broadly oblong, 1·5 mm long. Ovary globose, 1 mm long, white stellate-tomentose, 3-celled; style filiform, 3 mm long, with large white reflexed stellate hairs; ovules 2 per locule. Fruit a loculicidal capsule; seeds 1 per locule, almost cylindrical, villous, 2·5 mm long, 1 mm wide.

Distribution: South-west Western Australia, from Eneabba south to Gingin.

Eneabba Creek, C. A. Gardner 9128; 26½ mi from Watheroo on road to Jurien Bay, F. W. Humphreys 183; 5 mi SSE of Yeal Swamp in Wanneroo Forestry Reserve, Y. Chadwick 2554.

Lasiopetalum lineare has floral characteristics similar to those of *L. drummondii* Benth., but is distinguished by having linear leaves.

***Lasiopetalum rotundifolium* S. Paust sp. nov. (Figures 6 and 12)**
(rotundus = almost circular, folium = leaf)

Folia rotunda, profunde cordata, petiolata, 7–35 x 9–30 mm, supra rugosa glabra, subtus arcte stellato-tomentosa. *Cymae* pilis stellatis et simplici-moniliformibus tomentosae. *Bracteolae* solitariae, ovatae, ab calycibus distantes. *Calyx* 6 mm longus, intus glaber, extus pilis stellatis et simplici-moniliformibus tomentosus, lobis 5, ovato-lanceolatis. *Petala* nulla. *Antherae* fere sessiles, late ovato-oblongae, 2 mm longae. *Ovarium* 3-loculare, 1·5 mm longum, albo-stellato-tomentosum. *Stylus* 2 mm longus pilis stellatis grandibus reflexis albis hirsutus.

Type: 5–6 mi S of New Norcia, Western Australia, 1 Oct. 1947, C. A. Gardner 8690. Holo: PERTH, iso: CANB, K.

Erect, woody perennial, up to 40 cm high. Brauchlets tomentose with stellate and simple hairs. Leaves alternate; petiole 10–20 mm long, tomentose with stellate and simple-moniliform hairs; lamina orbicular, deeply cordate (the basal lobes sometimes imbricate), 7–35 mm long, 9–30 mm wide, rugose and glabrous above, closely grey stellate-tomentose below; margin ± crenulate slightly revolute. Cymes leaf-opposed, 20–40 mm long, ± 8-flowered, tomentose with stellate and simple-moniliform hairs; peduncle 10–20 mm long; bracts lanceolate, 3 mm long; bracteoles 1, ovate, 5 mm long, pink, papery, distant from calyx. Calyx pink (dark maroon near base of anthers), 6 mm long, almost glabrous within, tomentose with stellate and long simple-moniliform hairs outside, divided to near the base into 5 ovate lanceolate lobes. Petals absent. Anthers 5, almost-sessile, broadly ovate-oblong, dark maroon, 2 mm long. Ovary globose, 1·5 mm long, white stellate-tomentose, 3-celled; style simple, 2 mm long with large white reflexed stellate hairs; ovules 2 per locule. Seed not seen.

Distribution: South-west Western Australia; New Norcia area.

Swan River Colony, J. Drummond 119 (MEL 52366).

Lasiopetalum rotundifolium is known only from the Drummond and type collections. It has close affinities with *L. molle* Benth., but differs in having rounded, deeply cordate leaves.

***Lasiopetalum cardiophyllum* S. Paust. sp. nov. (Figures 7 and 12)**
(cardio = heart, phyllum = leaf)

Folia cordata, petiolata, 9–26 x 10–30 mm, coriacea, supra laevia glabra, subtus arcte stellata, tomentosa. *Cymae* pilis stellatis et simplici-moniliformibus tomentosae. *Bracteolae* solitariae, teretes, ab calycibus distantes. *Calyx* 8 mm longus, pilis stellatis et simplici-moniliformibus tomentosus, lobis 5, ovato-lanceolatis. *Petala* nulla. *Antherae* 5, fere sessiles, late ovato-oblongae, 1·5 mm longae. *Ovarium* 3-loculare, 1 mm longum, albo-stellato-tomentosum. *Stylus* 2·5 mm longus, pilis stellatis grandibus reflexis albis hirsutus.

Type: Mt. Saddleback, Darling Range, Western Australia, 13 Nov. 1904, A. Morrison s.n. Holo: PERTH.





Figure 6—*Lasiopetalum rotundifolium* sp. nov. Holotype—C. A. Gardner 8690.




Figure 7—*Lasiopetalum cardiophyllum* sp. nov. Holotype—Mt. Saddleback, A. Morrison.

Erect multistemmed, woody *perennial*, up to 40 cm high. *Branchlets* stellate-tomentose. *Leaves* alternate; petiole 4–15 mm long; lamina cordiform, 9–26 mm long, 10–30 mm wide, entire, coriaceous, smooth and almost glabrous above, closely grey stellate-tomentose below. *Cymes* leaf-opposed, 20–40 mm long, 5–many-flowered, sparsely tomentose with stellate and simple-moniliform hairs; peduncle 10–20 mm long; pedicels 10 mm long; bracts terete, 2 mm long, closely tomentose; bracteoles 1, terete, 2 mm long, closely tomentose, distant from calyx. *Calyx* pink (dark maroon near base of anthers), 8 mm long, tomentose with stellate and simple-moniliform hairs, divided to the base into 5 narrowly ovate-lanceolate acuminate lobes. *Petals* absent. *Anthers* 5, almost sessile, broadly ovate-oblong, 1.5 mm long, purple-black. *Ovary* globose, 1 mm long, white stellate-tomentose, 3-celled; style filiform, 2.5 mm long, with large white-reflexed stellate hairs; ovules 2 per locule. *Seed* not seen.

Distribution: South-west Western Australia; from North Bannister south to Mt. Saddleback.

Bunnings Plantation, 3 mi S of Halfway House on Albany Highway and 7 mi W of Highway, 21 Nov. 1969, R. J. Edmiston; towards Mt. Saddleback, between Williams and Hotham Rivers, 15 Nov. 1904, A. Morrison (K).

Lasiopetalum cardiophyllum has flowers similar to those of *L. floribundum* Benth. but is distinguished by having coriaceous, broadly ovate-cordate leaves which are stellate-tomentose below.

***Lasiopetalum glabratum* S. Paust. sp. nov. (Figures 8 and 12)**
(*glabratus* = glabrescent, referring to the leaves)

Folia late ovata ad semi-hastata vel cordata, petiolata, 10–30 x 6–30 mm, coriacea, fere glabra, supra reticulata, subtus manifeste nervata. *Cymae* pilis stellatis et simplici-moniliformibus tomentosae. *Bracteolae* solitariae, linearis, calyces subtus. *Calyx* 5 mm longus, sparsim stellato-tomentosus, extus etiam pilis simplici-moniliformibus, lobis 5, ovato-lanceolatis. *Petala* nulla. *Antherae* 5, fere sessiles, late ovato-oblongae, 1.5 mm longae. *Ovarium* 3-loculare, 1 mm longum, albo-stellato-tomentosum. *Stylus* 2.5 mm longus, pilis stellatis grandibus reflexis albis hirsutus.

Type: 2 mi N of turnoff to Mt. Cooke on Albany Highway, Western Australia, 6 Nov. 1971, Susan Paust 1068. *Holo:* PERTH.

Erect, multistemmed, woody *perennial*, up to 35 cm high. *Branchlets* stellate-tomentose. *Leaves* alternate; petiole stellate-tomentose, 3–6 mm long; lamina broadly ovate to semi-hastate, cordate, obtuse to acute, 10–30 mm long, 6–30 mm wide, coriaceous, undulate, almost glabrous, reticulate above, prominently veined below; margin semi-entire, slightly undulate and recurved. *Cyme* leaf-opposed, 30–50 mm long, 3–6-flowered, tomentose with stellate and simple-moniliform hairs; peduncle 20–30 mm long; pedicels 8 mm long; bracts linear, 2 mm long; bracteoles 1, linear, 2 mm long, subtending calyx. *Calyx* pink (dark maroon near base of anthers), 5 mm long, sparsely stellate-tomentose, the outer surface with additional simple-moniliform hairs, divided to the base into 5 ovate-lanceolate lobes. *Petals* absent. *Anthers* 5, almost sessile, broadly ovate-oblong, 1.5 mm long, purple-black. *Ovary* globose, 1 mm long, white stellate-tomentose, 3-celled; style filiform, 2.5 mm long, with large white reflexed stellate hairs; ovules 2 per locule. *Seed* not seen.

Distribution: South-west Western Australia: south-east of Perth, from Carmel east to York and south to Mt. Cooke.

Carmel, 29 Nov. 1928 E. Dell; 44 mi peg Albany Highway, A. R. Fairall 1711; between Perth and King George Sound, April, July 1854, W. H. Harvey; Swan River Colony, J. Drummond 20 (K).

This species has floral similarities to *L. floribundum* Benth. and *L. cardiophyllum* S Paust. (see above). It is distinguished from both in having coriaceous, undulate, almost glabrous leaves.




Figure 8—*Lasiopetalum glabratum* sp. nov. Holotype—S. Paust 1068.




Figure 9—*Lasiopetalum monticola* sp. nov. Holotype—Ellen Peak, A. Morrison.

Lasiopetalum monticola S. Paust. sp. nov. (Figures 9 and 12)

(monticola = a dweller on mountains)

Folia late ad anguste ovata, petiolata, 15–18 x 5–40 mm, supra laevia glabrescentia, subtus dense stellato-tomentosa. *Racemi* dense stellato-tomentosi. *Bracteolae* 3, filiformes, ab calyce distantes. *Calyx* 6 mm longus, intus sparsim stellato-hirsutus, extus dense albo-stellato-tomentosus, lobis 5, anguste ovatis. *Petala* 5, oblonga, 0.5 mm longa. *Antherae* 5, fere sessiles, anguste triangulares, 3 mm longae. *Ovarium* 3-loculare, 1 mm longum, papillatum. *Stylus* 2 mm longus, glaber.

Type: Ellen Peak, Stirling Range, Western Australia, 18 Oct. 1902, A. Morrison. Holo: PERTH, iso: CANB, K.

Erect, woody, *perennial* up to 50 cm high; branchlets, leaves and inflorescence densely stellate-tomentose, white to rufous. *Leaves* alternate; petiole 10–20 mm long; lamina broadly to narrowly ovate, entire, 15–80 mm long, 5–40 mm wide, smooth and becoming glabrous with impressed veins above, densely stellate-tomentose with prominent midrib below. *Raceme* leaf-opposed, 3–15-flowered, 40–110 mm long; peduncle 30–60 mm long; bracts terete, 5 mm long; bracteoles 3, filiform, 8 mm long, distant from calyx. *Calyx* 6 mm long, pink, sparsely white stellate-hairy within, densely white stellate-tomentose outside, divided almost to the base into 5 narrowly ovate lobes. *Petals* 5, oblong, concave, 0.5 mm long, glabrous. *Anthers* 5, almost sessile, narrowly triangular, 3 mm long. *Ovary* globose 1 mm long, papillate, 3-celled; style filiform, 2 mm long, glabrous, ovules 2 per locule. *Seed* not seen.

Distribution: South-west Western Australia; in the Stirling Range and on East Mt. Barren.

Stirling Range, E. Pritzel 701; Plantagenet Dist. Stirling Range, Diels 580; Mt. Warrungup, K. Newbey 1456; about half way along Salt River Road, Stirling Range, A. M. Ashby 1965; Eastern heights above Red Gum Springs, A. R. Fairall 465 and 471; 3 mi N of Ellen Peak, K. Newbey 315; East Mt. Barren, Sept. 1925. Gardner & Blackall; Gully, East Mt. Barren, Sept. 1924, A. Johnson.

Lasiopetalum monticola is similar in habit and appearance to *L. indutum* Steud. and *L. maxwellii* F. Muell., but is readily distinguished from these species by the papillate (rather than stellate-tomentose) ovary.

Lasiopetalum compactum S. Paust. sp. nov. (Figures 10 and 12)

(compactus = compact referring to the inflorescence)

Folia anguste oblonga, petiolata, 10–60 x 4–13 mm, coriacea, supra laevia glabra, subtus arcte stellato-tomentosa. *Cymae* in pseudo-capitula contractae. *Bracteolae* 3 lineares, calycem subtus. *Calyx* 5–8 mm longus, intus minute tuberculatus, extus dense albo-stellato-tomentosus, lobis 5, anguste ovatis acuminatis. *Petala* spatulata, 1 mm longa. *Stamina* 5, filamentis 1 mm longis, antheris anguste-triangularibus 2.5 mm longis. *Ovarium* 3-loculare, 2 mm longum, albo-stellato-tomentosum. *Stylus* 3 mm longus, glaber.

Type: 9 km SE of Ravensthorpe, Western Australia, 13 Aug. 1968, R. A. Saffrey 477. Holo: PERTH, iso: CANB, K.

Erect, woody, *perennial*, up to 1 m high. *Branchlets* closely stellate-tomentose, rufous to grey. *Leaves* alternate; petiole 5–7 mm long, closely stellate-tomentose; lamina narrowly oblong, entire, coriaceous, 10–60 mm long, 4–13 mm wide, smooth and glabrous with impressed midrib above, closely stellate-tomentose with prominent midrib below. *Cymes* leaf-opposed, 5–7-flowered, condensed into densely white stellate-tomentose heads; peduncle tomentose, 5–15 mm long; bracts linear, 5 mm long; bracteoles 3, linear, 7 mm long, subtending calyx. *Calyx* 5–8 mm long, pink and minutely tuberculate within, densely white stellate-tomentose outside, divided almost to the base into 5 narrowly ovate, acuminate lobes. *Petals* spatulate, slightly concave, 1 mm long glabrous. *Stamens* 5, filaments terete, 1 mm long; anthers narrowly triangular, 2.5 mm long. *Ovary* globose, 2 mm long, white stellate-tomentose, 3-celled; style filiform, 3 mm long, glabrous; ovules 2 per locule. *Fruit* a loculicidal capsule; seed 2 per locule, almost cylindrical, villous, 2 mm long, 1 mm wide.



HOLOTYPE

Western Australian Herbarium

Lasiopetalum compactum S. Moore

DETERMINAVIT Gerasa Aust. June 1973

WESTERN AUSTRALIAN HERBARIUM, PERTH
Flora of Western Australia

Loc.

Coll.

19




Figure 10—*Lasiopetalum compactum* sp. nov. Holotype—R. A. Saffrey 477.




Figure 11—Distribution of *Thomasia formosa* sp. nov., *T. gardneri* sp. nov., *T. microphylla* sp. nov. and *T. tremandroides* sp. nov.




Figure 12—Distribution of *Lasiopetalum cardiophyllum* sp. nov., *L. compactum* sp. nov., *L. glabratum* sp. nov., *L. lineare* sp. nov., *L. monticola* sp. nov., and *L. rotundifolium* sp. nov.

Distribution: South-west Western Australia, from the Fitzgerald River east to Mt. Burdett.

Fitzgerald R. valley, *R. D. Royce* 8906; Mt. Short, Ravensthorpe Range, *E. M. Bennett* 2494; 6 mi E of Ravensthorpe, *K. Newbey* 958; Between Ravensthorpe and Hopetoun, *S. Paust* 759; Yerritup Creek, 13 km N of coast at Stokes Inlet, *A. E. Orchard* 1188; Peak Charles, *C. Davies* 89; Mt. Burdett, 50 km NNE of Esperance, *E. N. S. Jackson* 1319; Swan River Colony, *J. Drummond* 264 (also MEL 52342, L.H. specimen).

The floral characters of *L. compactum* and *L. indutum* Steud. are similar and have previously caused confusion. *L. compactum*, however, is readily distinguished from the racemose *L. indutum* by the compact cyme and larger leaves.

New Synonyms in *Lasiopetalum*

1. *Lasiopetalum oppositifolium* F. Muell., *Fragm.* 2:5 (1860).

Type: Murchison River, Western Australia, *Oldfield*. Holo: MEL 52371.

L. angustifolium W. V. Fitzg., *J. West Austral. Nat. Hist. Soc.* No. 1:3 (1904), synon. nov.

Type: Geraldton, Western Australia, Sept. 1903, *W. V. Fitzgerald*. Iso: PERTH.

Lasiopetalum angustifolium was described by W. V. Fitzgerald as differing from *L. oppositifolium* by the presence of white reflexed hairs on the style. The examination of type and other herbarium collections has shown that the hairiness of the style is extremely variable, while the specimens are similar in other characters. It is therefore considered that the two are synonymous.

2. *Lasiopetalum indutum* Steud. in Lehmann, *Pl. Preiss.* 1:235 (1845).

Type: *Preiss* 1655. Iso: MEL 52353-4.

L. acutiflorum Turcz., *Bull. Soc. Imp. Naturalistes Moscou* 25:145 (1852), synon. nov.

Type: *J. Drummond* 254. Iso: MEL 52342, right hand specimen; MEL 52343; PERTH.

The two species are separated by Bentham in *Fl. Austral.* 1:259-260 (1863) by the degree of hairiness of the inside of the calyx lobes. From examination of the type and other herbarium collections, this character and leaf shape appear continuously variable. The two names are therefore considered synonymous.

Acknowledgement

Thanks are especially due to Mr. Alex George for the latin diagnoses.